

Amendments to the Claims:

- 1-5. (Cancelled)
6. (Currently Amended) A method for blocking apoptosis during preserving and storing a heart awaiting transplantation comprising:
perfusing said heart for up to 24 hours with a solution comprising:
- (a) a balanced isotonic solution in a physiologically acceptable amount;
 - (b) cyclosporin in an amount from about 2.5 μM to about 10 μM per liter of solution; and
 - (c) water.
7. (Original) The method according to Claim 6 wherein said balanced isotonic solution includes sodium, potassium, calcium, magnesium ions and bicarbonate.
8. (Cancelled)
9. (Original) The method according to Claim 6 wherein said cyclosporin is present in an amount from about 5.0 μM to about 8.0 μM per liter of solution.
10. (Currently Amended) The method according to Claim 6 wherein said balanced isotonic solution comprises:

Concentration Ranges in 1 Liter	
NaCl	85 mM to 145 mM
KCl	3 mM to 50 mM
CaCl ₂	0.5 mM to 2.5 mM
KH ₂ PO ₄	0.7 mM to 1.3 mM
MgSO ₄	0.9 mM to 4.8 mM
NaHCO ₃	15 mM to 35 mM
Glucose	1.0 mM to 50 mM

Appl. No.: 10/088,538

Filed: June 10, 2002

Page 3

~~and said cyclosporin is present in an amount from about 2.5 μ M to about 10 μ M per liter of solution.~~